




# Lesson 03: Functions




# Functions – an informal viewpoint




- Whenever one value “ $y$ ” depends on (or is determined by) another value “ $x$ ”, we say that  $y$  is a function of  $x$ . e.g.
  - Area  $A$  of a circle is a function of its radius  $r$ .
  - The temperature  $T$  at a place on a given day, is a function of time during the day ( $t$ ).
  - The height ( $H$ ) of a person is a function of the person’s age ( $x$ ).
- The function can be considered as a rule or a formula that gives the value  $y$ , when the value  $x$  is given to it.






# Function definition

- A **set** is a collection of objects. These objects are called **elements** of the set e.g.
  - $A = \{1, 4, 9, 16, 25, 36, 49\}$  or equivalently  $A = \{x: x = n^2, \text{ where } n \text{ is a natural number and } 1 \leq n \leq 7\}$
  - $B = \{a, e, i, o, u\}$
  - $C = \{x: 0 \leq x < \infty\}$
  - $D = [0, 1]$  (equivalent to  $\{x: 0 \leq x \leq 1\}$ )
  - $E = (0, 2]$  (equivalent to  $\{x: 0 < x \leq 2\}$ )
- A **function**  $f$  from set  $A$  to set  $B$  is a rule that assigns a *unique* (single) element  $f(x)$  in  $B$  to each element  $x$  in  $A$ . e.g.
  - $y = f(x) = x^2$ 
    - $f$  is the symbol used to denote the function.  $f(x)$  read as “ $f$  of  $x$ ” or the “value of  $f$  at  $x$ ” is the value that  $f$  assigns to any  $x$  in set  $A$ . Since  $f(x) = x^2$  here, the function  $f$  assigns the value  $x^2$  to each  $x$  in  $A$ .
    - $x$  and  $y$  are variables. Since  $y = f(x)$ ,  $y$  takes the value that the function  $f$  assigns to  $x$  in  $A$ .  $x$  can take on any value in  $A$  and the corresponding value of  $y$  depends on  $x$  (with the function  $f$  determining the value of  $y$  for a given  $x$ ). Therefore,  $x$  is called an **independent variable** while  $y$  is called a **dependent variable**.




# Function definition (continued)

- A function  $f$  from set  $A$  to set  $B$  is also denoted as  $f : A \rightarrow B$ . e.g.
  - $f : \mathbb{R} \rightarrow \mathbb{R}$  and  $y = f(x) = x^2$  ( $\mathbb{R}$  denotes the set of all real numbers).
- $A$  and  $B$  are usually sets of real numbers (like sets  $C$ ,  $D$  and  $E$  above).
- Function is denoted by letters like  $f$ ,  $g$ ,  $h$  etc.
- Set  $A$  is called the **domain** of the function. **When not stated explicitly, it is the largest set of real values  $x$  for which  $f(x)$  is real.**
- All values that  $f(x)$  can have (as  $x$  varies through the domain) is called the **range** of the function. Range is a subset of  $B$ .
- For  $y = f(x) = x^2$ , the domain is all real values  $x$   $(-\infty, \infty)$ , while the range is all real values  $\geq 0$   $[0, \infty)$ .
- Different values of  $x$  can be assigned to the same value by  $f$ . e.g.
  - $f(x_1) = f(x_2)$  for  $x_1 \neq x_2$  is a valid assignment by  $f$ .
- An **arrow diagram** has an arrow from  $x$  in set  $A$  to  $f(x)$  in set  $B$  to pictorially show the assignment done by the function. e.g. for  $f(x) = x^2$




# Function examples




- Area  $A$  of a circle is a function of its radius  $r$ 
  - $A = f(r)$ : This says that the variable  $A$  is the value of function  $f$  at  $r$ . But what is function  $f$ ?
  - $f(r) = \pi r^2$ : This says that the value of function  $f$  at  $r$  is equal to  $\pi r^2$ . Now the function is defined.
  - $A = f(r) = \pi r^2$ : This says that the variable  $A$  is equal to the value of the function  $f$  at  $r$ , which is equal to  $\pi r^2$ . Thus, we have expressed the circle area  $A$  as a function of radius  $r$ .
- Consider  $f(x) = 3x^2 + 8$ 
  - $f(1) = 3.(1)^2 + 8 = 11$
  - $f(2) = 3.(2)^2 + 8 = 20$
  - Domain of the function is all real values  $x$ .
  - Range is  $[8, \infty)$ . Why?  $x^2 \geq 0$ , so the min value of  $3x^2 + 8$  is 8.


# Some functions and their graphs



**Graph of a function**  $f(x)$  is the set of all points  $(x, y)$  in the coordinate plane such that  $y = f(x)$ .  $x$  takes all values in the domain of  $f(x)$ .




Odd Powers of  $x$   
Note range for these functions is  $(-\infty, \infty)$



Even Powers of  $x$   
Note range for these functions (except the constant function) is  $[0, \infty)$


Polynomial function of degree  $n$  has the form  $f(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$  ( $a_n \neq 0$ )

# Common functions (continued)



Greatest integer function written as  $y = f(x) = \lfloor x \rfloor$  assigns to each  $x$ , the greatest integer less than or equal to  $x$ .

Note the range consists of all integers. Also  $y$  jumps from one integer to the next, e.g. from 2 to 3, as  $x$  changes from "a little less than 3" to 3.



The graph consists of the blue lines and the dots.


The sloping black line  $y = x$ , acts as a reference.

Dots (found at integer values) indicate that the point belongs to the graph of  $\lfloor x \rfloor$ .


Absolute value function  $y = f(x) = |x|$  is defined as

$$y = x, \text{ when } x \geq 0$$

$$= -x \text{ when } x < 0$$



# Sum, difference, product and quotient of functions



For every  $x$  in the domain of both functions  $f(x)$  and  $g(x)$ ; the sum, difference, product and quotient are defined.

$$\text{Sum } (f + g)(x) = f(x) + g(x)$$

e.g. the value of the sum function  $(f + g)$  at  $x$ , is the value of the function  $f$  at  $x$ , plus the value of the function  $g$  at  $x$ .

$$\text{Difference } (f - g)(x) = f(x) - g(x)$$

$$\text{Product } (fg)(x) = f(x)g(x)$$

$$\text{Quotient } \left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)} \quad \text{Also, } g(x) \text{ must not be } 0$$

Example:

$$\text{Let } f(x) = \sqrt{x+1} \text{ and } g(x) = \sqrt{4-x^2}$$

Domain of  $f(x)$  is  $[-1, \infty)$  (since  $x + 1 \geq 0$ )

Domain of  $g(x)$  is  $[-2, 2]$  (since  $4 - x^2 \geq 0$ )

Points common to both domains  $[-1, 2]$

Therefore,

$$(f + g)(x) = \sqrt{x+1} + \sqrt{4-x^2} \text{ with domain } [-1, 2]$$

$$(f - g)(x) = \sqrt{x+1} - \sqrt{4-x^2} \text{ with domain } [-1, 2]$$


$$(fg)(x) = \sqrt{(x+1)(4-x^2)} \text{ with domain } [-1, 2]$$

$$\left(\frac{f}{g}\right)(x) = \sqrt{\frac{x+1}{4-x^2}} \text{ with domain } [-1, 2] \text{ since } 4 - x^2 \text{ must not be } 0.$$

$$\left(\frac{g}{f}\right)(x) = \sqrt{\frac{4-x^2}{x+1}} \text{ with domain } (-1, 2] \text{ since } x+1 \text{ must not be } 0$$



# Composite functions



If  $f(x)$  and  $g(x)$  are functions, then the composite function  $f \circ g$  (“ $f$  composed with  $g$ ”) is defined as  $(f \circ g)(x) = f(g(x))$  [the value of function  $f$  at  $g(x)$ ]

- Given a  $x$  in the domain of  $g$ 
  - function  $g$  assigns (maps) it to  $g(x)$ .
  - If  $g(x)$  lies in the domain of  $f$ , then  $f(x)$  assigns the value  $f(g(x))$  to it.
- Therefore, composite  $(f \circ g)$  is defined for all  $x$  where  $g(x)$  lies in the domain of  $f$ : basically all  $x$ , for which both  $g(x)$  and  $f(g(x))$  are defined.
- $(g \circ f)(x) = g(f(x)) \neq (f \circ g)(x) = f(g(x))$

Example

Let  $f(x) = \sqrt{x+3}$  and  $g(x) = 2x$

$(f \circ g)(x) = f(g(x)) = f(2x) = \sqrt{2x+3}$  with domain  $[-3/2, \infty)$

$(g \circ f)(x) = g(f(x)) = g(\sqrt{x+3}) = 2\sqrt{x+3}$  with domain  $[-3, \infty)$

$(f \circ f)(x) = f(f(x)) = f(\sqrt{x+3}) = \sqrt{\sqrt{x+3}+3}$

Since  $\sqrt{x+3} \geq 0$  for  $x \in [-3, \infty)$ , the domain is  $[-3, \infty)$

$(g \circ g)(x) = g(g(x)) = g(2x) = 4x$  with domain  $(-\infty, \infty)$

